

NUTRACEUTICAL VALUE OF THE NEW STRAWBERRY CULTIVARS

Jadwiga Żebrowska, Jacek Gawroński, Elżbieta Kaczmarska,
Magdalena Dyduch-Siemińska, Izabella Jackowska,
Marzena Pabich

University of Life Sciences in Lublin

Abstract. Regarding an increasing interest of consumers towards berry fruit, as they have been proved to have potential health benefits, the nutraceutical fruit value of seven strawberry cultivars: 'Felina', 'Feltar', 'Hulta', 'Jota', 'Pastel', 'Plena' and 'Teresa' selected at the University of Life Sciences in Lublin, Plant Genetics and Breeding Department was evaluated in this study. In fruits harvested at maturity, the level of phytochemical compounds such as sugars, vitamin C, and polyphenols including quercetin and gallic acid as well as antiradical activity was estimated. Among all genotypes tested, the cultivar 'Jota' showed the highest content of glucose and fructose as well a high content of sucrose, therefore the total sugar content exceeded 12 g per 100 g of fresh weight (FW). Similar parameters in terms of content of total sugars and the individual fractions have cultivar 'Pastel'. Cultivars with low sucrose content were 'Feltar' and 'Teresa', but as regards monosaccharide's cvs. 'Plena' and 'Felina'. The content of vitamin C in fruits of the cultivars tested showed significant differences. It ranged from 64.5 mg·100 g⁻¹ FW in cv. 'Jota' to 104.33 mg·100 g⁻¹ FW in cv. 'Plena'. The highest level of antiradical activity occurred in cultivars 'Jota', 'Teresa' and 'Feltar' (90.1%, 87.1% and 86.3% DPPH; respectively). The results obtained indicated that the differentiation of the chemical composition of strawberry fruit is clearly dependent upon the genetic diversity of cultivars tested, what we argue that all of them grown under the same climatic and soil conditions using the same agricultural technology.

Key words: antioxidant activity, DPPH, polyphenols, fruit quality, sugars, vitamin C

INTRODUCTION

The cultivated strawberry *Fragaria × ananassa* Duch. is the most widely distributed small fruit crop in the world. It is grown in every country with a temperate or subtropi-

Corresponding author: Jacek Gawroński, Department of Genetics and Horticultural Plant Breeding, University of Life Sciences in Lublin, Akademicka 15, 20-950 Lublin, e-mail: jacek.gawronski@up.lublin.pl

cal climate and even in many tropical countries in highland areas, where the climate is mild. Breeding of the species was initiated with selection of hybrids of *F. chiloensis* × *F. virginiana* and hybrids obtained from crosses with other wild representatives of *Fragaria* genus. The breeding was undertaken by amateurs and private firms in England in the XIX c. to obtain plants producing big fruits [Darrow 1966, Staudt 1989]. Since the middle of the XX c., strawberry breeders have been focused on improving the plant's resistance to diseases, productivity and local adaptation, as well as on improving fruit quality [Hancock et al. 2008].

In recent years, as a result of the increasing requirements of food consumers, greater attention has been placed on the nutraceutical value of fruits. Nutraceutical, a portmanteau of the words “nutrition” and “pharmaceutical”, is a food or food product that reportedly provides health and medical benefits, including the prevention and treatment of disease. The term “nutraceutical” was originally defined by Dr. Stephen L. DeFelice, founder and chairman of the Foundation of Innovation Medicine (FIM), Crawford, New Jersey [www.fimdefelice.org 2011]. The evidence-based health benefits of berry consumption provide a means of promoting strawberries [Schöppllein et al. 2002, Cordenunsi et al. 2003]. Strawberry fruits are a valuable component of the diet, since they are of low caloric content and contain easily assimilable sugars, as well as several bioactive phytochemicals, including vitamin C and other antioxidant compounds such as polyphenols in that phenolic acids and flavonoids. Phenolic classes commonly found in strawberries are hydroxybenzoic acids (gallic and ellagic acids), hydroxycinnamic acids (p-cumaric). Flavonoids are grouped in several structural classes including anthocyanins (pelargonidin-3-glycoside pigment), flavones, flavan-3-ols (catequins), flavanones, flavonols (quercetin, kaempferol and myricetin) and hydrolysable tannins (ellagitanins). *In vivo* and *in vitro* studies have been recently carried out to establish the effects of strawberries on free radical scavenging and on the prevention of cardiovascular diseases [Mateos et al. 2005, Pinto et al. 2008].

In 1970s, at the Genetics and Horticultural Plant Breeding Department of the Faculty of Horticulture (nowadays Faculty of Horticulture and Landscape Architecture), University of Life Sciences in Lublin, the strawberry breeding program had been undertaken to select new cultivars of this species. This program was carried out to improve the fruit yield and its quality i.e. the fruit weight, shape, texture, colour, taste and firmness. As a result of this breeding work based on the conventional clone selection, some new cultivars were released in that: ‘Felina’, ‘Feltar’, ‘Hulta’, ‘Jota’, ‘Pastel’, ‘Plena’ and ‘Teresa’. All of them originated from the hybrid seedlings obtained after crosses between different parental cultivars of the species. Regarding an increasing interest of consumers towards berry fruit, as they have been proved to have potential health benefits, the nutraceutical fruit value of these seven cultivars selected at the University of Life Sciences in Lublin was evaluated in this study.

MATERIAL AND METHODS

The field experiment was established in August 2009 in triplicates with 20 plants per each cultivar at the Felin Experimental Station (N 51°13', E 22°39'). During the experiment typical agronomic procedures recommended for strawberry plantations [Żurawicz et al. 2005] were used. The fruit samples were evaluated in the first and second year of

full yielding of plants i.e. in 2011 and 2012. Samples of 500 g of full-ripened strawberry fruit of the cultivars 'Felina', 'Feltar', 'Hulta', 'Jota', 'Pastel', 'Plena' and 'Teresa' (pedigree of cultivars is presented in Table 1.) were hand-harvested twice in June 2011 and 2012. After harvest the fruit samples were delivered to the laboratory where they were manually washed and carefully sorted to remove any damaged ones. The level of phytochemical compounds such as sugars (glucose, fructose and sucrose), vitamin C, total polyphenols including flavonoids and phenolics as well as antiradical activity was measured in fruits.

Table 1. Pedigree of cultivars investigated

Cultivar	Pedigree
Felina	Ottawa × Purpuratka
Feltar	(Senga Tigaiga × Merton Dawn)S ₁
Hulta	Senga Litessa × Torrey
Jota	(Tioga S ₁ × Cambridge Vigour S ₁) × Dukat
Pastel	Paula × (Senga Sengana × Talisman)
Plena	Senga Sengana × Merton Dawn
Teresa	Redgauntlet S ₁ × Senga Sengana S ₁

Evaluation of vitamin C and soluble sugars content

Determination of the vitamin C and soluble sugars (glucose, fructose and sucrose) content in fruits of cultivars tested was conducted at the Central Laboratory of Agricoology, University of Life Sciences in Lublin with the use of the HPLC method in accordance with PN-EN 12630:2002 and PN-EN 14130:2003 standards (respectively).

Polyphenols assay

Fruit extract preparation. Fruit extract was prepared from 10 g of fresh samples using 40 mL of 80% (by volume) aqueous ethanol. The mixture was extracted (in water bath at 80°C), kept for 20 min in inert atmosphere, and filtered through a Whatman filter paper using a Büchner funnel. Extraction of the residue was repeated under the same conditions. The filtrates were combined and diluted to 100 mL in volumetric flask with 80% aqueous ethanol, and the obtained extract was used for determination of total polyphenols (phenolics, flavonoids) as well as DPPH assay.

Phenolics content of fruit extracts was measured by Folin-Ciocalteu's phenol reagent [Kim et al. 2003]. First, 200 µl of appropriately diluted sample or gallic acid standard were added to 2.6 ml of distilled deionized water. Then, 200 µl of Folin-Ciocalteu's phenol reagent was added at time zero and mixed. After 6 min, 2 ml of 7% (w/v) Na₂CO₃ solution was added and mixed. Absorbance was measured after incubation for 90 min at room temperature; absorbance was measured at 750 nm versus a prepared blank. The blank consisted of 200 µl 50% (v/v) methanol instead of sample. Gallic acid in 50% (v/v) methanol solution in concentrations of 10, 30, 60 and 100 mg·l⁻¹ was used as a standard and a calibration curve was drawn for each day of analysis. The content of phenolics was expressed as mg gallic acid equivalent (GAE)·100 g⁻¹ of fresh weight. All samples were analyzed in five replicates.

Flavonoids content of fruit extracts was determined using a spectrophotometric method [Floegel et al. 2010]. 500 µl of sample or standard (quercetin) were mixed with 3.2 ml of distilled deionized water. At time zero, 150 µl of 5% (w/v) sodium nitrite

(NaNO₂) solution were added and mixed. After 5 min, 150 µl of 10% (w/v) aluminum chloride (AlCl₃) was added and mixed. After 6 min, 1 ml of 1 M sodium hydroxide (NaOH) was added and mixed. Absorbance of the colored flavonoid-aluminum complex was measured immediately at 510 nm versus a blank. The blank consisted of 500 µl of 50% (v/v) methanol instead of sample. For standard quercetin was dissolved in 50% (v/v) methanol to concentrations of 10, 30, 60 and 100 mg·l⁻¹. A calibration curve was drawn. Flavonoids content in the strawberry extract was expressed as mg quercetin equivalent (QE)·100 g⁻¹ fresh sample. All samples were analyzed in five replicates.

Evaluation of antiradical activity (DPPH)

Antioxidant activity was determined by the DPPH (2, 2-diphenyl-1-picrylhydrazyl) radical – scavenging method developed by Brand-Williams et al. [1995] slightly modified by Kim et al. [2002]. DPPH radical solution was prepared by dissolving 0.02 g of DPPH in 50 ml of methanol (analytical grade, POCh). From this solution 5 ml was taken and diluted with methanol to a volume of 50 ml. In this way the final concentration of the radical 0.04 mg/ml was obtained. To 100 ml of the extract was added 3 ml of DPPH (a concentration of 0.04 mg/ml). Samples were mixed; the tube was capped with a stopper and allowed to stand for 30 minutes. After this time the absorbance was measured at a wavelength of 517 nm. A control was prepared by diluting 100 µl methanol (analytical grade) in 3 ml of DPPH. Antioxidant activity was calculated according to the formula: $AA = [1 - (A_p / A_k)] \cdot 100\%$, where: AA – antioxidant activity, A_p – absorbance of the sample, A_k – absorbance of the control sample. Antioxidant activity was expressed as the percent of inhibition. Results obtained underwent statistical analysis with the use SAS Enterprise Guide 3.0 and Excel 3.0 statistical software. The significance of differences between means was established by the Tukey test at $p \leq 0.05$.

RESULTS AND DISCUSSION

Sugars. The results indicated the existence of a varietal diversity in relation to the content of glucose, fructose and sucrose in the fruit of the cultivars tested (tab. 2). Among them cv. 'Jota' showed the highest content of glucose and fructose and also a high sucrose therefore as the only one exhibited a total sugar content exceeding 12 g per 100 g FW. Cultivar with similar parameters in terms of total sugar and individual fractions was 'Pastel'. The group of varieties of low sucrose content included 'Feltar' and 'Teresa', while monosaccharide's 'Plena' and 'Felina'. In the Koszański et al. [2006] studies cultivar 'Senga Sengana' had the highest content of total sugars. However, their content was lower as compared to the cultivars 'Pastel' and 'Jota', but similar to the 'Plena' cultivar. In other studies of these authors the cultivars 'Elsanta' and 'Elkat' had lower sugar content, even compared to the lowest in this respect cultivars investigated in this work. In contrast Rochalska et al. [2011] obtained a very high sugar content (in range of 28.93–30.98 g·100 g⁻¹ FW) significantly exceeding these both values presented in this work and in works of other authors [Koszański et al. 2005, Kopytowski et al. 2006, Mishra et al. 2015]. Content of reducing sugars obtained for all cultivars tested were similar to the results obtained by Skupień [2003] for 'Pandora', 'El-

santa', 'Senga Sengana', 'Vicoda', 'Marmolada' and 'Kent' strawberry cultivars, however, the sucrose content compared to the cultivars studied by this author was significantly higher. Impact on the level of these characteristics other than cultivar has climatic, agronomic and edaphic factors [Koszański et al. 2006].

Table 2. The sugars content ($\text{mg} \cdot 100 \text{ g}^{-1}$ FW) in the fruits of cultivars tested

Cultivar	Sugars			Total sugar
	sucrose	glucose	fructose	
Felina	4644 a*	2932 cd	3007 b	10583 a
Feltar	3439 c	3259 bc	3655 ab	10353 a
Hulta	4171 abc	3137 bc	3605 ab	10913 a
Jota	4416 ab	4166 a	4152 a	12701 a
Pastel	4325 ab	3631 b	3670 ab	11626 a
Plena	4151 abc	2588 d	2889 b	9628 a
Teresa	3692 bc	3179 bc	3409 ab	10280 a

* – means followed by the same letter are not significantly different in columns at $p \leq 0.05$

Vitamin C. The vitamin C content in the fruit of the cultivars tested showed significant differences (tab. 3). The lowest occurred in the cultivar 'Jota', while the highest in cv. 'Plena'. The value of this trait in studies of other authors also reached a high degree of volatility and amounted to $47.5 \text{ mg} \cdot 100 \text{ g}^{-1}$ FW in 'Senga Sengana', [Koszański et al. 2006], $81.27 \text{ mg} \cdot 100 \text{ g}^{-1}$ FW in 'Selva' [Mortazavi et al. 2014]. However, in studies conducted by Mishra et al. [2015] the value of this trait ranged from 62.67 to $73.6 \text{ mg} \cdot 100 \text{ g}^{-1}$ FW (on average for twenty genotypes), Nunes and Delgado [2014] reported values in the range of 60.1–71.8 $\text{mg} \cdot 100 \text{ g}^{-1}$ FW while in studies by Lal et al. [2013] varied between 51.03 and $89.40 \text{ mg} \cdot 100 \text{ g}^{-1}$ FW. It was also observed, that genotypes with lower content of vitamin C (tab. 3) had higher contents of sugars (tab. 2). This was in line with Mishra et al. [2015] reports about statistically negative correlation between these properties at both genotypic (-0.906) and phenotypic (-0.773) level. However, as Aubert and Bosc [2014] stated changes in the content of vitamin C in fruit distribution are irrelevant (although some think otherwise – Mortazavi et al. 2014), but significantly reduces the level of sucrose.

Flavonoids. Differentiation of flavonoids content expressed as quercetin equivalent ($\text{mg QE} \cdot 100 \text{ g}^{-1}$ FW) in the fruits of cultivars studied proved to be highly significant and ranged from 192.40 to $707.41 \text{ mg QE} \cdot 100 \text{ g}^{-1}$ FW (tab. 3). The best in this regard the variety 'Jota' ($707.41 \text{ mg QE} \cdot 100 \text{ g}^{-1}$ FW) included almost 4× more of this compound than 'Teresa'. Cheel et al. [2007] determined the flavonoid content between 30 mg quercetin equivalents $\cdot 100 \text{ g}^{-1}$ fresh fruit for the white strawberry to 123.2 mg quercetin equivalents $\cdot 100 \text{ g}^{-1}$ fresh fruit for *F × ananassa* cv. 'Chandler', while Meyers et al. [2003] reported the highest flavonoid content for cultivars 'Evangeline' and 'Earliglow' and the lowest for 'Sparkle' and 'Allstar'. Wang and Lewers [2007] indicated significant differences in the content of flavonoids between the cultivars of strawberries, which were reflected in the results of this work. Crespo et al. [2010] recognized the levels of the properties of a significant feature of the cultivar mainly conditioned by the action of genetic factors, and only slightly environmental factors. However Andreotti et

al. [2014], noticed differences in the content of this compound in the fruit of cultivar ‘Elsanta’ from plantations with different habitat conditions (especially altitude). It must therefore be regarded as Aubert and Bosc [2014] suggested, that the level of the characteristics analyzed was a cultivar’s feature, wherein one must take into account the fact that the fruits which are not fully mature contain lower concentration of this compound in compared to the ripe fruit [Mahmood et al. 2012].

Table 3. The content of antioxidants in fruits of cultivars investigated and their antiradical activity

Cultivar	Vitamin C (mg·100 g ⁻¹ FW)	Polyphenols compounds		Polyphenols content (mg·100 g ⁻¹ FW)	Antiradical activity DPPH (% of inhibition)
		flavonoids (mg QE·100 g ⁻¹ FW)	phenolics (mg GAE·100 g ⁻¹ FW)		
Felina	88.76 abc*	344.51 c	321.25 bc	665.76 bc	80.08 d
Feltar	74.85 cd	222.58 ef	416.18 a	638.76 bc	86.30 b
Hulta	79.35 bcd	249.82 e	408.39 a	658.22 bc	84.38 bc
Jota	64.50 d	707.41 a	335.95 abc	1043.36 a	90.06 a
Pastel	71.66 d	484.70 b	305.63 c	790.33 b	82.18 cd
Plena	104.33 a	295.45 d	318.40 c	613.85 c	75.73 e
Teresa	90.92 ab	192.40 f	400.46 ab	592.86 c	87.07 ab

* – means followed by the same letter are not significantly different in columns at $p \leq 0.05$

Phenolics. Phenolics content expressed as gallic acid equivalent varied from 305.63 in cultivar ‘Pastel’ to 416.18 mg GAE in ‘Feltar’ (tab. 3). In the study of Pineli et al. [2012] phenolic acids content ranged from 180 to 280 mg GAE·100 g⁻¹ FW, in Lal et al. [2013] ranged from 380.1 (cv. ‘Douglas’) to 888.1 (cv. ‘Howard’) mg GAE·100 g⁻¹ FW, while in Wang et al. [2014] noted 210 mg GAE·100 g⁻¹ FW. The differences in the content of these compounds between strawberry cultivars were also observed by Proteggente et al. [2002], Scalzo et al. [2005], Djilas et al. [2011], while between 62 fruit species by Fu et al. [2011]. Among the strawberry cultivars tested the highest concentration of phenolic compounds with antioxidant activity was characterized by the cultivar ‘Jota’ (tab. 3). Significantly lower level was observed in the cultivar ‘Pastel’, while the lowest occurred in cv. ‘Teresa’. Different levels of this trait between genotypes were also observed by Scalzo et al. [2005] and Debnath and Ricard [2009]. Wang and Lewers [2007] showed that the fruit of *Fragaria virginiana* accessions had significantly higher antioxidant capacity and total phenolics content (expressed as GAE) than the fruit from different *Fragaria* × *ananassa* cultivars. Capocasa et al. [2008] stated that limited knowledge is available on the possibility of improving strawberry nutritional traits by breeding; however they showed an improvement in fruit nutritional quality in the breeding material originated from *F. virginiana* ssp. *glauca* inter-specific crosses. According to Khanizadeh et al. [2014], despite the fact that the content of phenolic compounds and antioxidant properties depended mainly on genetic factors, however, the production system can significantly affect at their level. The providing of fruit nutritional and nutraceutical quality can be achieved by appropriate post-harvest fruit treatment [Wang et al. 2014].

DPPH. The present data provided the evidence that strawberry genotypes tested differed in their activity to react and quench DPPH radicals (tab. 3). Cultivar ‘Jota’ was the

best for an antioxidant activity, while ‘Plena’ was the worst. Cheel et al. [2007] accessed DPPH activity for cultivated strawberry at the level of 86% whereas native accessions characterized lower activity (42%). Previous studies conducted by other researchers reported the variation in antioxidant activity among 10 strawberry cultivars and 9 advances selections [Debnath and Ricard 2009]. As a reason for variation of this trait Anttonen et al. [2006] indicated a fruit order in inflorescence and planting date, whereas Wang and Zheng [2001] saw it in a day/night temperature but above all growing variety. Floegel et al. [2011] classified strawberries among species with a high antioxidant capacity.

Table 4. The correlation and determination coefficients matrix between antiradical activity (DPPH) and antioxidant components content for 7 strawberry cultivars

Cultivar	Antioxidant components	
	flavonoids (QE)	phenolics (GAE)
Felina	0.78 ¹	-0.27
	0.61 ²	0.07
Feltar	0.29	-0.26
	0.08	0.07
Hulta	-0.46	0.32
	0.21	0.10
Jota	0.45	-0.45
	0.20	0.20
Pastel	0.95*	0.42
	0.90	0.18
Plena	0.41	0.04
	0.17	0.002
Teresa	0.98*	0.34
	0.97	0.12
Mean	0.49	0.22
	0.24	0.05

¹ – correlation coefficient, ² – determination coefficient, * – correlation significant at $p \leq 0.05$

Interrelations between the antiradical activity (DPPH) and antioxidant compounds content in fruits of cultivars tested are presented in Table 4, but only a few of them were statistically significant (at cultivar level). A high positive correlation between antiradical activity and flavonoids (QE) content for cultivars ‘Teresa’ and ‘Pastel’ indicated that this compound could be one of the main components responsible for antioxidant activity of these fruits. The rest of cultivars demonstrated the diversity in this respect that was especially evident in ‘Hulta’ cultivar for which the coefficient of correlation had negative value. In respect to the influence of phenolics (GAE) on the antioxidant capacity it should be regarded that this dependence was the most diverse in this study. For cvs. ‘Pastel’, ‘Teresa’ and ‘Hulta’ the impact of phenolics on the antioxidant activity was positive and amounted to 0.42, 0.34 and 0.32 respectively, while for the cvs. ‘Jota’, ‘Felina’ and ‘Feltar’ was negative. Only in cultivar ‘Hulta’ the antiradical activity of phenolics (GAE) was higher than of flavonoids. Nevertheless in this group of cultivars the comparison of antiradical activity between flavinoid and phenolic compounds showed that flavonoids (QE) determined the capacity for DPPH radicals scavenging

almost 5× more than phenolics (GAE) (tab. 4). So far in the literature there is a scarce information about correlation between the antioxidant activity and antioxidant compounds content at the level of single cultivar. In Scalzo et al. [2005] study total antioxidant capacity and phenolics content were correlated negatively (for cultivars ‘Don’, ‘Idea’, ‘Camarosa’, ‘Onda’, ‘Sveva’, ‘Patty’), whereas in Pineli et al. [2012] research for cultivars ‘Osogrande’ and ‘Camino Real’ vice versa. Positive correlations between DPPH activity and phenolic content were also observed in Meyers et al. [2003], Cheel et al. [2007], Djilas et al. [2011] and Huang et al. [2012] research. Also strong positive relationship between DPPH and phenolics (Spearman-Rho coefficients = 0.897) and flavonoids (0.708) content were obtained by Floegel et al. [2011]. Wang and Lin [2000] reported a significant correlation between antioxidant capacity and both total phenolics and vitamin C, and estimated on the basis the high values of a determination coefficient ($R^2 = 0.8093$, $R^2 = 0.812$; respectively). In this study, the high values of R^2 (tab. 4) were obtained only in the case of cultivar ‘Pastel’ and ‘Teresa’ for flavonoids (QE) content. This means that especially in the case of the last cultivar antioxidant capacity resulted from the content of flavonoids (fig. 1). Therefore, as stated Almeida et al. [2011] and Meyers et al. [2003] the antioxidant activities in the fruits cannot be attributed solely to their phenolics contents, but also to the actions of different antioxidant compounds present in the fruits which may act additively and synergistically with others and final activity may be dependent upon relative proportions of each or presence/absence of particular compounds.




Fig. 1. Relationship between antiradical activity (DPPH) and flavonoids content in cultivar ‘Teresa’

CONCLUSIONS

In our study the differences in the bioactive compounds content as well as in the antiradical activity (with DPPH) observed between cultivars resulted mainly from their genotypic diversity, since all cultivars grew under the same climatic and soil conditions using the same agricultural technology. The highest nutraceutical value showed the fruits of cv. 'Jota' due to the soluble sugars content and the highest value of antiradical activity. In this group of cultivars the antioxidant activity of bioactive compounds was mainly determined by flavonoids expressed as quercetin equivalent. In general, on the basis of the present results it should be stated that the nutraceutical fruit value depended not only on the bioactive phytochemicals content and relative proportions between them but also on the interrelated actions of different chemical compounds present in the fruits. This study also illustrated the existence of wide ranges of variations among strawberry cultivars that provides opportunities for genetic gain through hybridization and selection. Regarding an importance of the knowledge on the strawberry nutraceutical properties for consumers' health, further studies in this area aimed at the estimation of these characteristics individually for each cultivar are needed.

REFERENCES

- Almeida, M.M.B., de Suosa, P.H.M., Arriaga, A.M.C., do Prado, G.M., Magalhaes, C.E.C., Maia, G.A., de Lemos, T.L.G. (2011). Bioactive compounds and antioxidant activity of fresh exotic fruits from northeastern Brazil. *Food Res. Int.*, 44, 2155–2159.
- Andreotti, C., Guerrero Chavez, G., Zago, M. (2014). Quality of strawberry fruit cultivated in a highland area in South Tyrol (Italy): First results. *Acta Hort.*, 1049, 795–799.
- Anttonen, M.J., Hoppula, K.I., Nestby, R., Verheul, M.J., Karjalainen, R.O. (2006). Influence of fertilization, mulch color, early forcing, fruit order, planting date, growing environment and genotype on the content of selected phenolics in strawberry (*Fragaria* × *ananassa* Duch.) fruits. *J. Agric. Food Chem.*, 54, 2614–2620.
- Aubert, C., Bosc, J.P. (2014). Postharvest evolution of aromatic and nutritional quality for various strawberry cultivars in France. *Acta Hort.*, 1049, 789–794.
- Brand-Williams, W., Cuvelier, M.E., Berset, C. (1995). Use of a free radical method to evaluate antioxidant activity LWT. *Food Sci. Tech.*, 28, 25–30.
- Capocasa, F., Scalzo, J., Mezzetti, B., Battino, M. (2008). Combining quality and antioxidant attributes in the strawberry, the role of genotype. *Food Chem.*, 111(4), 872–878.
- Cheel, J., Theoduloz, C., Rodriguez, J.A., Caligari, P.D.S., Schmeda-Hirschmann, G. (2007). Free radical scavenging activity and phenolic content in achnes and thalamus from *Fragaria chiloensis* ssp. *chiloensis*, *F. vesca* and *F. × ananassa* cv. Chandler. *Food Chem.*, 102, 36–44.
- Cordenunsi, B.R., Nascimento, J.R.O., Lajolo, F.M. (2003). Physico-chemical changes related to quality of five strawberry fruit cultivars during cool-storage. *Food Chem.*, 83, 167–173.
- Crespo, P., Bordonaba, J., Terry, L., Carlen, C. (2010). Characterization of major taste and health – related compounds of four strawberry genotypes grown at different Swiss production sites. *Food Chem.*, 122, 16–24.
- Darrow, G.M. (1966). The strawberry – history, breeding and physiology. The New England Institute for Medical Research, <http://www.nal.usda.gov/pgdic/strawberry/booksevt.htm>.pp.21

- Debnath, S.C., Ricard, E. (2009). ISSR, anthocyanin content and antioxidant capacity analyses to characterize strawberry genotypes. *J. App. Hort.*, 11(2), 83–89.
- Djilas, S.M., Tepić, A.N., Savatović, S.M., Šumić, Z.M., Čanadanović-Brunet, J.M., Četković, G.S., Vulić, J.J. (2011). Chemical composition and antioxidant activity of two strawberry cultivars. *Acta Period. Tech.*, 42, 33–44.
- Floegel, A., Kim, D.O., Chung, S.J., Song, W.O., Fernandez, M.L., Bruno, R.S., Koo, S.I., Chun, O.K. (2010). Development and validation of an algorithm to establish a total antioxidant capacity database of the US diet. *Int. J. Food Sci. Nutr.*, 61, 600–623.
- Floegel, A., Kim, D.O., Chung, S.J., Koo, S.I., Chun, O.K. (2011). Comparison of ABTS/DPPH assay to measure antioxidant capacity in popular antioxidant-rich US food. *J. Food Comp. Anal.*, 24, 1043–1048.
- Fu, L., Xu, B.T., Xu, X.R., Gan, R.Y., Zhang, Y., Xia, E.Q., Li, H.B. (2011). Antioxidant capacities and total phenolic content of 62 fruits. *Food Chem.*, 129, 345–350.
- Hancock, J.F., Sjulín, T.M., Lobos, G.A. (2008). Strawberries. In: *Temperate fruit crop breeding*, Hancock J.F. (ed.). Springer Science + Business Media B.V., 93–438.
- Huang, W., Zhang, H., Liu, W., Li, C. (2012). Survey of antioxidant capacity and phenolic composition of blueberry, blackberry, and strawberry in Nanjing. *J. Zhejiang Univ. Sci., B.*, 13(2), 94–102.
- Khanzadeh, S., Fan, L., Fang, Ch., Charles, M.T., Shutian, T. (2014). Effect of production systems on phenolic composition of strawberry fruits. *Acta Hort.*, 1049, 513–516.
- Kim, D.O., Chun, O.K., Kim, Y.J., Moon, H.Y., Lee, C.Y. (2003). Quantification of polyphenolics and their antioxidant capacity in fresh plums. *J. Agric. Food Chem.*, 51, 6509–6515.
- Kim, D.O., Lee, K.W., Lee, H.J., Lee, C.Y. (2002). Vitamin C equivalent antioxidant capacity (VCEAC) of phenolic phytochemicals. *J. Agric. Food Chem.*, 50, 3713–3717.
- Kopytowski, J., Kawecki, Z., Bojarska, J.E., Stany, V. (2006). Ocena plonowania i jakości owoców kilku odmian truskawki uprawianej na Warmii. *Zesz. Nauk. Inst. Sadow. Kwiac.*, 14, 53–60.
- Koszański, Z., Rumas-Rudnicka, E., Karczmarczyk, S., Rychter, P. (2005). Wpływ nawadniania i nawożenia mineralnego na plonowanie i cechy jakościowe dwóch odmian truskawek uprawianych na glebie lekkiej. *Inż. Roln.*, 3(63), 251–256.
- Koszański, Z., Rumas-Rudnicka, E., Podsiadło, C. (2006). Wpływ nawadniania kropowego i nawożenia mineralnego na jakość owoców truskawki. *J. Elementol.*, 11(1), 21–27.
- Lal, S., Ahmed, N., Singh, S.R., Shing, D.B., Sharma, O.C., Kumar, R. (2013). Variability of health and bioactive compounds in strawberry (*Fragaria* × *ananassa* Duch.) cultivars grown under an Indian temperate ecosystem. *Fruits*, 68, 423–434.
- Mahmood, T., Anwar, F., Abbas, M., Saari, N. (2012). Effect of maturity on phenolics (phenolic acids and flavonoids) profile of strawberry cultivars and mulberry species from Pakistan. *Int. J. Mol. Sci.*, 13, 4591–4607; doi:10.3390/ijms13044591
- Mateos, R., Lecumberri, H., Ramos, S., Goya, L., Brawo, L. (2005). Determination of malondialdehyde (MDA) by high-performance liquid chromatography in serum and liver as a biomarker for hypercholesterolemia and evaluation of the effect of diets rich in phenolic antioxidants from fruits. *J. Chromat., B.*, 827, 76–82.
- Meyers, K.J., Watkins, C.B., Pritts, M.P., Liu, R.H. (2003). Antioxidant and antiproliferative activities of strawberries. *J. Agr. Food Chem.*, 51, 6887–6892.
- Mishra, P.K., Ram, R.B., Kumar, N. (2015). Genetic variability, heritability and genetic advance in strawberry (*Fragaria* × *ananassa* Duch.). *Turk. J. Agric. For.*, 39, doi:10.3906/tar-1408-99

- Mortazavi, S.M.H., Siruie, B., Moalemi, N., Eshghi, S. (2014). The effects of polyamines and UV-C irradiation on postharvest quality of strawberry fruit. *Acta Hort.*, 1049, 749–754.
- Nunes, M.C.N., Delgado, A. (2014). Quality of organic compared to conventionally grown strawberries at the retail level. *Acta Hort.*, 1049, 723–730.
- Pineli, L., Moretti, C.L., Chiarello, M.D. (2012). Quality, bioactive compounds and antioxidant activity of strawberries grown in Brazilian savannah and stored at different temperatures. *J. Food Agric. Environ.*, 10(2), 165–171.
- Pinto, M.S., Kwon, Y., Apostolidis, E., Lajolo, F.M., Genovese, M.I., Shetty, K. (2008). Functionality of bioactive compounds in Brazilian strawberry (*Fragaria* × *ananassa* Duch.) cultivars: Evolution of hyperglycemia and hypertension potential using *in vitro* models. *J. Agr. Food Chem.*, 56, 4386–4392.
- PN-EN 12630:2002 (2002). Soki owocowe i warzywne. Oznaczanie zawartości glukozy, fruktozy, sorbitolu i sacharozy. Metoda wysokosprawnej chromatografii cieczowej. Sektor Żywności Rolnictwa i Leśnictwa, Organ Techniczny – KT 38 ds. Przetworów Owocowych i Warzywnych. ICS 67.160.20.
- PN-EN 14130:2003 (2003). Artykuły Żywnościowe – Oznaczanie witaminy C za pomocą HPLC. Sektor Żywności Rolnictwa i Leśnictwa, Organ Techniczny – KT 235 ds. Analizy żywności. ICS 67.050.
- Proteggente, A.R., Sekher-Pannala, A., Paganga, G.A., Van Buren, L., Wagner, E., Wiseman, S. (2002). The antioxidant activity of regular consumed fruit and vegetables reflects their phenolic and vitamin C composition. *Free Rad. Res.*, 36, 217–233.
- Rochalska, M., Orzeszko-Rywka, A., Czaplą, K. (2011). The content of nutritive substances in strawberries according to cropping system. *J. Res. Appl. Agric. Engin.*, 56(4), 84–86.
- Scalzo, J., Politi, A., Mezzetti, B., Battino, M. (2005). Plant genotype affects total antioxidant capacity and phenolic content in fruit. *Nutrition*, 21, 207–213.
- Schöpplein, E., Krüger, E., Rechner, A., Hoberg, E. (2002). Analytical and sensory qualities of strawberry cultivars. *Acta Hort.*, 567, 805–808.
- Skupień, K. (2003). Ocena wybranych cech jakościowych świeżych i mrożonych owoców sześciu odmian truskawki. *Acta Sci. Pol. Hortorum Cultus*, 2(2), 115–123.
- Staudt, G.S. (1989). The species of *Fragaria*, their taxonomy and geographical distribution. *Acta Hort.*, 265, 23–34.
- Wang, S.Y., Lewers, K. (2007). Antioxidant capacity and flavonoid contents in wild strawberries. *J. Amer. Soc. Hort. Sci.*, 132(5), 647–658.
- Wang, B.G., Feng, X.Y., Li, W.S., Yang, J.J., Yang, Y., Shi, L. (2014). Effect of a Ma box combined with high carbon dioxide on preservation qualities of strawberry. *Acta Hort.*, 1049, 743–747.
- Wang, S.Y., Lin, H.S. (2000). Antioxidant activity in fruits and leaves of blackberry, raspberry and strawberry varies with cultivar and development stage. *J. Agr. Food Chem.*, 48, 140–146.
- Wang, S.Y., Zheng, W. (2001). Effect of plant growth temperature on antioxidant capacity in strawberry. *J. Agr. Food Chem.*, 49(10), 4977–4982.
- www.fimdefelice.org/p2466.html. The NutraCeutical Revolution: Fueling a Powerful, New International Market". *Fimdefelice.org*. [retrieved: 2011-06-03].
- Żurawicz, E., Bielenin, A., Lisek, J., Łapanowska, B.H., Machecki, J., Treder, W. (2005). Metodyka integrowanej produkcji truskawek. Plantpress. Kraków.

NUTRACEUTYCZNA WARTOŚĆ NOWYCH ODMIAN TRUSKAWKI

Streszczenie. Biorąc pod uwagę wzrastające zainteresowanie konsumentów spożyciem owoców jagodowych ze względu na ich udowodniony korzystny wpływ na ludzkie zdrowie, w obecnych badaniach oceniano nutraceutyczne właściwości owoców siedmiu odmian truskawki: 'Felina', 'Feltar', 'Hulta', 'Jota', 'Pastel', 'Plena' i 'Teresa' wyselekcjonowanych na Uniwersytecie Przyrodniczym w Lublinie, w Katedrze Genetyki i Hodowli Roślin Ogrodniczych. W owocach zebranych w fazie pełnej dojrzałości oznaczono zawartość związków fitochemicznych, takich jak cukry, witamina C i polifenole ogółem oraz oszacowano aktywność antyoksydacyjną (z DPPH). Spośród badanych odmian, owoce odmiany 'Jota' charakteryzowały się największą zawartością glukozy i fruktozy, a także sacharozy – całkowita zawartość cukrów przekroczyła $12 \text{ g} \cdot 100 \text{ g}^{-1}$ świeżej masy (f.w.). Podobne wartości pod względem całkowitej zawartości cukrów i ich poszczególnych frakcji zawierały owoce odmiany 'Pastel'. W owocach odmian 'Feltar' i 'Teresa' stwierdzono najmniejszą zawartość sacharozy, natomiast w owocach odmian 'Pleny' i 'Feliny' – najmniejszą ilość cukrów prostych. Zawartość witaminy C w owocach badanych odmianach była bardzo zróżnicowana i wynosiła od $64.5 \text{ mg} \cdot 100 \text{ g}^{-1}$ f.w. ('Jota') do $104.33 \text{ mg} \cdot 100 \text{ g}^{-1}$ f.w. ('Plena'). Odmiany 'Jota', 'Teresa' i 'Feltar' wykazywały najwyższą zdolność do znoszenia wolnych rodników (odpowiednio 90.1, 87.1 i 86.3% DPPH). Wszystkie analizowane odmiany były uprawiane w tych samych warunkach klimatyczno-glebowych i poddawane takim samym zabiegom agrotechnicznym, stąd też na podstawie uzyskanych wyników wnioskuje się, że skład chemiczny owoców truskawki wyraźnie zależy od zróżnicowania genetycznego badanych genotypów.

Słowa kluczowe: aktywność antyoksydacyjna, DPPH, jakość owoców, polifenole, cukry, witamina C

Accepted for print: 15.02.2016

For citation: Żebrowska, J., Gawroński, J., Kaczmarska, E., Dyduch-Siemińska, M., Jackowska, I., Pabich, M. (2016). Nutraceutical value of the new strawberry cultivars. *Acta Sci. Pol. Hortorum Cultus*, 15(3) 71–82.